arXiv:1210.4142v2 [quant-ph] 18 Oct 2012

Quantum Theory Contains Maximally Incompatible Observables

Paul Busch[*]
Department of Mathematics, University of York, York, United Kingdom

Teiko Heinosaari[†] and Jussi Schultz[‡]
Turku Centre for Quantum Physics, Department of Physics and Astronomy, University of Turku, FI-20014 Turku, Finland

We show that Quantum Theory contains observables that are as incompatible as any probabilistic physical theory can have. We define the joint measurability region of a probabilistic theory which describes the amount of added noise needed to make any pair of observables in the theory jointly measurable. We then show that the joint measurability region of Quantum Theory is contained in the joint measurability region of any other probabilistic theory.

PACS numbers: 03.65.Ta

Quantum Theory has a number of important features not known in classical physics, ranging from the superposition and indeterminacy principles formulated by the pioneers to the more recently discovered no-cloning and no-broadcasting theorems. It is an old problem to identify operationally significant properties of Quantum Theory that distinguish it from other probabilistic theories. In recent years many features have been under intensive investigation from this perspective, including information processing [1], optimal state discrimination [2], entropy [3], purification [4] and discord [5]. It has been found that some properties are quite generally valid in any nonclassical (no-signaling) probabilistic theories while others are specifically quantum.

The existence of pairs of incompatible observables marks one of the most striking distinctions between Quantum Theory and classical physical theories. There are many manifestations of incompatibility, perhaps the most famous being the Heisenberg Uncertainty Principle [6]. In this Letter we demonstrate that Quantum Theory contains observables that are as incompatible (in a sense to be defined) as observables in any probabilistic theory can be.

Our aim is thus to compare Quantum Theory to other possible probabilistic physical theories. We first need to set some minimal constraints. A probabilistic theory is a framework that provides a description of physical systems in terms of states and observables with the following general properties:

(i) The states of a system are represented by the elements of a convex subset of a real vector space.

(ii) An observable is represented as an affine mapping from the set of states into the set of probability distributions on some outcome space. For simplicity, we restrict ourselves here to observables with a finite or countable number of outcomes.

(iii) Any affine mapping from the set of states into the set of probability distributions is a valid observable.

We consider a particular probabilistic theory (PT) as given by a family of convex sets of states with associated sets of observables that share some properties specific to that PT. One may think of each pair consisting of a set of states with associated set of observables as an *instance* of a PT representing a particular type of physical system.

Given a PT, we denote by $\mathfrak{M}(j|\varrho)$ the probability of obtaining a measurement outcome j when an observable $\mathfrak{M}$ is measured in a state ϱ. Hence, $0 \leq \mathfrak{M}(j|\varrho) \leq 1$ and $\sum_j \mathfrak{M}(j|\varrho) = 1$. We will typically label the measurement outcomes by integers.

In Quantum Theory the states are described by density operators and observables correspond to POVMs [7]. Their duality is given by the trace formula (with ϱ a density operator and M a POVM)

$$\mathfrak{M}(j|\varrho) = \mathrm{tr}\left[\varrho \mathsf{M}(j)\right] . \qquad (1)$$

Another example of a probabilistic theory is a classical theory, where the states are probability measures on a phase space Ω and observables are traditionally represented as functions $m : \Omega \to \mathbb{R}$; the associated affine maps from states ϱ to probability distributions are then given by the formula

$$\mathfrak{M}(j|\varrho) = \varrho(\{x \in m^{-1}(j)\}) . \qquad (2)$$

Continuing our discussion on general probabilistic theories, we note that it follows from the required properties (i)-(iii) that the set of observables is a convex set; a mixture of two observables is an observable. Physically mixing corresponds to an experiment where we switch between two measurement apparatuses with a random probability. We can directly write a mixture of two observables with the same set of measurement outcomes. If the sets of measurement outcomes differ, we can still write a mixture by first adding enough outcomes and then embedding both sets into $\mathbb{Z}$.

Another consequence of the basic requirements is that every constant mapping $\varrho \mapsto p$, where p is a fixed probability distribution, is an observable and we call it a *trivial*

---

[*] Electronic address: pb516@york.ac.uk
[†] Electronic address: teiko.heinosaari@utu.fi
[‡] Electronic address: jussi.schultz@utu.fi

observable. A trivial observable $\mathfrak{T}$ corresponds to a dice rolling experiment, where we randomly pick the measurement outcome according to a given fixed probability distribution, without manipulating the state at all. In Quantum Theory, trivial observables are described by POVMs T such that each operator $\mathsf{T}(j)$ is a multiple of the identity operator, i.e., $\mathsf{T}(j) = t_j I$ for some $0 \leq t_j \leq 1$ with $\sum_j t_j = 1$.

The concept of *joint measurement* can be defined in any probabilistic theory. Two observables $\mathfrak{M}_1$ and $\mathfrak{M}_2$ are *jointly measurable* if there exists an observable $\mathfrak{M}$ such that

$$\sum_k \mathfrak{M}(j,k|\varrho) = \mathfrak{M}_1(j|\varrho)\,, \sum_j \mathfrak{M}(j,k|\varrho) = \mathfrak{M}_2(k|\varrho)\,. \tag{3}$$

In this case $\mathfrak{M}$ is called a *joint observable* of $\mathfrak{M}_1$ and $\mathfrak{M}_2$. If $\mathfrak{M}_1$ and $\mathfrak{M}_2$ are not jointly measurable, then we say that they are *incompatible.*

Any probabilistic theory contains jointly measurable pairs of observables. Namely, a trivial observable $\varrho \mapsto p$ is jointly measurable with any other observable; we can write a joint observable

$$\mathfrak{M}(j,k|\varrho) = \mathfrak{M}_1(j|\varrho)p(k) \tag{4}$$

for the trivial observable and any other observable $\mathfrak{M}_1$. This simply corresponds to an experiment where we measure $\mathfrak{M}_1$ and simultaneously roll a dice. It is a well known fact that, in Quantum Theory, an observable which is jointly measurable with all other observables is necessarily a trivial observable. Indeed, any POVM element of such an observable commutes with all projections and must therefore be a scalar multiple of the identity (e.g. [8, Theorem IV.1.3.1]).

The following simple observation is a key ingredient for our discussion.

Proposition 1. *Let $\mathfrak{M}_1$ and $\mathfrak{M}_2$ be two observables and $0 \leq \lambda \leq 1$. Then $\lambda\mathfrak{M}_1 + (1-\lambda)\mathfrak{T}_1$ and $(1-\lambda)\mathfrak{M}_2 + \lambda\mathfrak{T}_2$ are jointly measurable for any choice of trivial observables $\mathfrak{T}_1$ and $\mathfrak{T}_2$.*

This proposition can be proved with the following construction. First, let p_1 and p_2 be the probability distributions related to $\mathfrak{T}_1$ and $\mathfrak{T}_2$. We define an observable $\mathfrak{M}$ by formula

$$\mathfrak{M}(j,k|\varrho) = \lambda p_2(k)\, \mathfrak{M}_1(j|\varrho) + (1-\lambda)p_1(j)\, \mathfrak{M}_2(k|\varrho)\,. \tag{5}$$

For a fixed ϱ, the right hand side is clearly a probability distribution. Moreover, the right hand side is an affine mapping on ϱ; therefore $\mathfrak{M}$ is an observable. The marginal observables are

$$\sum_k \mathfrak{M}(j,k|\varrho) = \lambda\mathfrak{M}_1(j|\varrho) + (1-\lambda)p_1(j),$$
$$\sum_j \mathfrak{M}(j,k|\varrho) = (1-\lambda)\mathfrak{M}_2(k|\varrho) + \lambda p_2(k).$$

This proves Prop. 1.



FIG. 1: (Color online) The region $J(\mathsf{M}_x, \mathsf{M}_y)$ for two orthogonal spin-$\frac{1}{2}$ measurements is a quadrant of the unit disk. The region $\triangle$ (light) is a subset of $J(\mathfrak{M}_1, \mathfrak{M}_2)$ for any pair $\mathfrak{M}_1, \mathfrak{M}_2$, while the surplus region (dark) depends on the specific pair under consideration.

The physical idea behind this construction is the following. In each measurement run we flip a coin and, depending on the result, we measure either $\mathfrak{M}_1$ or $\mathfrak{M}_2$ in the input state ϱ. In this way we get a measurement outcome for either $\mathfrak{M}_1$ or $\mathfrak{M}_2$. In addition to this, we roll a dice and pretend that this is a measurement outcome for the other observable. In this way we get an outcome for both observables simultaneously. The overall observable is the one given in formula (5).

For two observables $\mathfrak{M}_1$ and $\mathfrak{M}_2$, we denote by $J(\mathfrak{M}_1, \mathfrak{M}_2)$ the set of all points $(\lambda, \mu) \in [0,1] \times [0,1]$ for which there exist trivial observables $\mathfrak{T}_1, \mathfrak{T}_2$ such that $\lambda\mathfrak{M}_1 + (1-\lambda)\mathfrak{T}_1$ and $\mu\mathfrak{M}_2 + (1-\mu)\mathfrak{T}_2$ are jointly measurable, and we call $J(\mathfrak{M}_1, \mathfrak{M}_2)$ the *joint measurability region* of $\mathfrak{M}_1$ and $\mathfrak{M}_2$. The joint measurability region thus characterizes how much noise (in terms of trivial observables) we need to add to be able to have a joint realization of $\mathfrak{M}_1$ and $\mathfrak{M}_2$. Clearly, $\mathfrak{M}_1$ and $\mathfrak{M}_2$ are jointly measurable if and only if $(1,1) \in J(\mathfrak{M}_1, \mathfrak{M}_2)$.

The joint measurability region $J(\mathfrak{M}_1, \mathfrak{M}_2)$ is a convex region which can be plotted in the plane. This follows as a simple application of Prop. 2 of [9]; with the help of this lemma it also follows from Prop. 1 that

$$\triangle \equiv \{(\lambda,\mu) \in [0,1] \times [0,1] : \lambda + \mu \leq 1\} \subseteq J(\mathfrak{M}_1, \mathfrak{M}_2)\,.$$

As an example, suppose that we are within Quantum Theory and $\mathfrak{M}_1$ and $\mathfrak{M}_2$ correspond to spin-$\frac{1}{2}$ measurements in two orthogonal directions, say x and y -axes. We then describe them with two POVMs M_x and M_y, where

$$\mathsf{M}_x(\pm 1) = \tfrac{1}{2}(I \pm \sigma_x)\,, \quad \mathsf{M}_y(\pm 1) = \tfrac{1}{2}(I \pm \sigma_y)\,, \tag{6}$$

and σ_x, σ_y are the usual Pauli matrices in $\mathbb{C}^2$. It has been shown in [10] that for the uniformly distributed trivial observable $\pm 1 \mapsto \frac{1}{2}I$ (hence describing an unbiased coin), the two observables $\lambda\mathsf{M}_x + (1-\lambda)\frac{1}{2}1$ and $\mu\mathsf{M}_y + (1-\mu)\frac{1}{2}1$ are jointly measurable if and only if $\lambda^2 + \mu^2 \leq 1$. By Prop. 3 of [9] this inequality is a necessary condition for the joint measurability of any pair $\lambda\mathsf{M}_x + (1-\lambda)\mathsf{T}_1$ and $\mu\mathsf{M}_y + (1-\mu)\mathsf{T}_2$, where $\mathsf{T}_1, \mathsf{T}_2$ are arbitrary trivial observables. Therefore, we conclude that

$$J(\mathsf{M}_x, \mathsf{M}_y) = \{(\lambda,\mu) \in [0,1] \times [0,1] : \lambda^2 + \mu^2 \leq 1\}\,. \tag{7}$$



(a) (b)

FIG. 2: (Color online) The region $\triangle$ (light) is a subset of the joint measurability region J_{PT} (colored) in any probabilistic theory. The larger the surplus region (dark) is, the more jointly measurable the theory globally is. If (a) and (b) would be joint measurability regions for two different probabilistic theories, then we would conclude that (b) is more incompatible than (a).

This region is depicted in Fig. 1.

We are now ready to define the main concept of this Letter. A global joint measurability feature of a probabilistic theory PT is characterized by the intersection of all the sets $J(\mathfrak{M}_1, \mathfrak{M}_2)$ across all instances of PT, and we denote

$$\begin{aligned} J_{PT} =\{&(\lambda,\mu) \in [0,1]\times[0,1] : (\lambda,\mu) \in J(\mathfrak{M}_1,\mathfrak{M}_2) \\ &\text{for all pairs of observables } \mathfrak{M}_1 \text{ and } \mathfrak{M}_2 \\ &\text{in all instances of PT}\}. \end{aligned}$$

We call J_{PT} the *joint measurability region* of PT. We always have $\triangle \subseteq J_{PT}$, but J_{PT} can be larger than $\triangle$. The larger the surplus region is, the more jointly measurable the theory is globally; see Fig. 2. If $(\lambda,\mu) \notin J_{PT}$, this means that there is a pair of observables $\mathfrak{M}_1$ and $\mathfrak{M}_2$ such that the mixtures $\lambda\mathfrak{M}_1 + (1-\lambda)\mathfrak{T}_1$ and $\mu\mathfrak{M}_2 + (1-\mu)\mathfrak{T}_2$ are incompatible with any choice of trivial observables $\mathfrak{T}_1$ and $\mathfrak{T}_2$.

Since J_{PT} can be defined in any probabilistic theory, we can compare the joint measurability regions of different probabilistic theories. We obviously have $J_{PT} = [0,1]\times[0,1]$ in any probabilistic theory where all measurements are jointly measurable, such as the classical probability theory. In the most incompatible case we would have $J_{PT} = \triangle$. We will next show that Quantum Theory is, globally, as incompatible as a probabilistic theory can be.

Theorem 1. *In Quantum Theory $J_{QT} = \triangle$. In particular, $J_{QT} \subseteq J_{PT}$ for any probabilistic theory PT.*

In Quantum Theory every observable $\mathfrak{M}$ corresponds to a unique POVM M by equation (1). We will prove that for any pair $(\lambda,\mu) \notin \triangle$, there are quantum observables M_1 and M_2 such that the mixtures $\lambda\mathsf{M}_1 + (1-\lambda)\mathsf{T}_1$ and $\mu\mathsf{M}_2 + (1-\mu)\mathsf{T}_2$ are incompatible with any choice of trivial observables $\mathsf{T}_1, \mathsf{T}_2$. Our proof is based on a recent result [11] on the joint measurability region for two complementary observables, which is a generalization of the result illustrated in Fig. 1.

Proof of Theorem 1. We have earlier seen that $\triangle \subseteq J_{PT}$, so we need to show that $J_{QT} \subseteq \triangle$. Let $(\lambda,\mu) \notin \triangle$, i.e., $\lambda+\mu > 1$. Fix $\epsilon > 0$ such that $\lambda+\mu > 1+\epsilon$. We then choose d to be a positive integer satisfying

$$\frac{\sqrt{d}-1}{d-1} \leq \epsilon\,. \tag{8}$$

(This can be done since the left hand side → 0 when $d \to \infty$.) We will consider a quantum system that is described by a d-dimensional Hilbert space $\mathcal{H}_d$. Let $\{\varphi_j\}_{j=0}^{d-1}$ be an orthonormal basis for $\mathcal{H}_d$. We define another orthonormal basis $\{\psi_k\}_{k=0}^{d-1}$ for $\mathcal{H}_d$ by

$$\psi_k = 1/\sqrt{d}\sum_j e^{-2\pi i\frac{jk}{d}}\varphi_j\,. \tag{9}$$

The orthonormal bases $\{\varphi_j\}_{j=0}^{d-1}$ and $\{\psi_k\}_{k=0}^{d-1}$ are mutually unbiased, i.e., $|\langle\,\varphi_j\,|\,\psi_k\,\rangle| = \textit{constant}$. We define two POVMs M_1 and M_2 by

$$\mathsf{M}_1(j) = |\varphi_j\rangle\langle\varphi_j|\,,\quad \mathsf{M}_2(k) = |\psi_k\rangle\langle\psi_k|\,. \tag{10}$$

We thus obtain a pair of d-outcome observables on $\mathcal{H}_d$. Since M_1 and M_2 consist of projections and $\mathsf{M}_1(j)\mathsf{M}_2(k) \neq \mathsf{M}_2(k)\mathsf{M}_1(j)$, it follows that they are incompatible.

As proved in [11], the observables $\lambda'\mathsf{M}_1 + (1-\lambda')\mathsf{T}_1$ and $\mu'\mathsf{M}_2 + (1-\mu')\mathsf{T}_2$ are incompatible for any choice of trivial observables $\mathsf{T}_1, \mathsf{T}_2$ whenever

$$\lambda' + \mu' > 1 + \frac{\sqrt{d}-1}{d-1}\,. \tag{11}$$

Since

$$\lambda+\mu > 1+\epsilon \geq 1 + \frac{\sqrt{d}-1}{d-1}\,, \tag{12}$$

we conclude that $(\lambda,\mu) \notin J_{QT}$. $\square$

In the proof of Theorem 1 we have used quantum observables with arbitrarily many (but a finite number of) outcomes. It is natural to ask if this is a necessary trick in order to reach the conclusion $J_{QT} = \triangle$. To present a partial answer to this question, let us investigate the joint measurability region in the case of pairs of binary quantum observables. Our aim is to show that

$$\{(\lambda,\mu) \in [0,1]\times[0,1] : \lambda^2+\mu^2 \leq 1\} \subseteq J(\mathsf{M}_1,\mathsf{M}_2)$$

for any binary observables M_1 and M_2, regardless of the dimension of the Hilbert space. In other words, we will show that two orthogonal spin observables are as incompatible as any binary observables can be.

To this end, let us note that two binary quantum observables are incompatible if and only if they enable a violation of the Bell-CHSH inequality [12]. We must therefore look at the Bell expression

$$\mathcal{B} = |\langle\mathsf{M}_1\mathsf{N}_1\rangle + \langle\mathsf{M}_1\mathsf{N}_2\rangle + \langle\mathsf{M}_2\mathsf{N}_1\rangle - \langle\mathsf{M}_2\mathsf{N}_2\rangle|\,.$$



(a) (b)

FIG. 3: In (a) the grey area represents the possible values that α and β can obtain by varying the observables and the state in the Bell expression $\mathcal{B} = |\alpha + \beta|$. The solid lines represent the Tsirelson bound $\mathcal{B} = 2\sqrt{2}$ and the dashed lines represent the bound $\mathcal{B} = 2$. By considering only observables which are mixtures with the uniformly distributed trivial observable with fixed λ and μ, the area becomes smaller as depicted in (b), and a suitable choice of weights makes the violation of the Bell-CHSH inequality impossible.

Let us denote $\alpha = \langle \mathsf{M}_1\mathsf{N}_1\rangle + \langle \mathsf{M}_1\mathsf{N}_2\rangle$ and $\beta = \langle \mathsf{M}_2\mathsf{N}_1\rangle - \langle \mathsf{M}_2\mathsf{N}_2\rangle$. Using, e.g., [13, Theorem 1] one verifies that by varying the observables M_j and N_j and the bipartite state the numbers α and β obtain all of the values from the disc $\alpha^2 + \beta^2 \leq 4$. If we now mix the observables M_j with the trivial observable $\mathsf{T}(\pm 1) = \frac{1}{2}I$ with some weights λ and μ we see that the pair (α, β) turns into $(\lambda\alpha, \mu\beta)$, thus changing the Bell expression from $|\alpha + \beta|$ to $|\lambda\alpha + \mu\beta|$. We must therefore determine those (λ, μ) for which $|\lambda\alpha + \mu\beta| \leq 2$ for all (α, β) satisfying $\alpha^2 + \beta^2 \leq 4$ (see Fig. 3). But the boundary curve for this region is obtained when the equations $(\alpha/\lambda)^2 + (\beta/\mu)^2 = 4$ and $\alpha + \beta = 2$ have at most one common solution. By inserting $\beta = 2 - \alpha$ into the first equation the problem reduces to determining when the discriminant is negative or zero, and one readily verifies that this is the case exactly when $\lambda^2 + \mu^2 \leq 1$.

In conclusion, given any pair of binary observables M_1 and M_2, and weights λ and μ with $\lambda^2 + \mu^2 \leq 1$, the mixtures $\lambda\mathsf{M}_1 + (1-\lambda)\mathsf{T}$ and $\mu\mathsf{M}_2 + (1-\mu)\mathsf{T}$ can not be used to violate the Bell-CSHS inequality and must therefore be jointly measurable. We note that in the case $\mu = \lambda$ the same result using a different technique has been obtained by Banik *et al.* [14].

We note that in the proof of Theorem 1 we have used quantum observables acting on an arbitrarily high (but finite) dimensional (but finite) Hilbert space. We can again ask if this is a necessary trick in order to reach the conclusion $J_{QT} = \triangle$. Since two mutually unbiased bases are expected to be among the most incompatible observables in a fixed dimension d, our construction suggests that the conclusion $J_{QT} = \triangle$ can be reached only if one considers arbitrarily high dimensions. A proof of this fact is, however, lacking.

We show finally that the conclusion $J_{QT} = \triangle$ can be reached by using a *single pair* of incompatible observables if we consider an infinite dimensional system and observables with infinite number of outcomes.

Let $\mathcal{H}$ be an infinite dimensional Hilbert space and write it as a direct sum of finite d-dimensional Hilbert spaces $\mathcal{H}_d$, $\mathcal{H} = \bigoplus_{d=2}^{\infty} \mathcal{H}_d$. In each $\mathcal{H}_d$ consider a pair of mutually unbiased orthonormal bases $\{\varphi_j^d\}_{j=0}^{d-1}$ and $\{\psi_k^d\}_{k=0}^{d-1}$, where the latter is obtained from the first one by the formula (9). We define two POVMs N_1 and N_2 via

$$\mathsf{N}_1(d,j) = |\varphi_j^d\rangle\langle\varphi_j^d| \,, \quad \mathsf{N}_2(d,k) = |\psi_k^d\rangle\langle\psi_k^d| \,. \tag{13}$$

These observables act in the infinite dimensional Hilbert space $\mathcal{H}$ and d in (13) is an index labeling the different outcomes. The outcome space of N_1 and N_2 is $\Omega_\infty \equiv \{(d,j) : d = 2, 3, \ldots, j = 0, \ldots, d-1\}$.

Theorem 2. *The observables* N_1 *and* N_2 *defined in* (13) *satisfy* $J(\mathsf{N}_1, \mathsf{N}_2) = \triangle$.

Proof of Theorem 2. Let p_1 and p_2 be two probability distributions defined on Ω_∞. Assume that $\lambda + \mu > 1$ and define two observables $\mathsf{N}_{1,\lambda}, \mathsf{N}_{2,\mu}$ via

$$\begin{aligned} \mathsf{N}_{1,\lambda}(d,j) &= \lambda\, |\varphi_j^d\rangle\langle\varphi_j^d| + (1-\lambda)\, p_1(d,j) I, \\ \mathsf{N}_{2,\mu}(d,k) &= \mu\, |\psi_k^d\rangle\langle\psi_k^d| + (1-\mu)\, p_2(d,k) I \,. \end{aligned} \tag{14}$$

We need to show that $\mathsf{N}_{1,\lambda}$ and $\mathsf{N}_{2,\mu}$ are incompatible. To prove this, we make a counter assumption that $\mathsf{N}_{1,\lambda}, \mathsf{N}_{2,\mu}$ are jointly measurable. This implies that for any projection P on $\mathcal{H}$, the projected observables $P\mathsf{N}_{1,\lambda}P$ and $P\mathsf{N}_{2,\mu}P$ acting on a subspace $P\mathcal{H}$ are jointly measurable. (If G is a joint observable of two observables $\mathsf{M}_1, \mathsf{M}_2$, then $P\mathsf{G}P$ is a joint observable of $P\mathsf{M}_1P, P\mathsf{M}_2P$ in $P\mathcal{H}$.) Especially, the projections of $\mathsf{N}_{1,\lambda}$ and $\mathsf{N}_{2,\mu}$ to any subspace $\mathcal{H}_d$ should be jointly measurable. But from the result cited in the proof of Theorem 1 we know that for d large enough, the projections to $\mathcal{H}_d$ are incompatible. Hence, $\mathsf{N}_{1,\lambda}$ and $\mathsf{N}_{2,\mu}$ are incompatible. □

We note that the observables N_1 and N_2 defined in (13) are not the only pair that satisfy $J(\mathsf{N}_1, \mathsf{N}_2) = \triangle$. Namely, we can modify N_1 and N_2 in any chosen subspace $\mathcal{H}_d$ but the conclusion $J(\mathsf{N}_1, \mathsf{N}_2) = \triangle$ is still true since it depends on the fact that N_1 and N_2 contain mutually unbiased bases in arbitrarily high dimension. An interesting question within Quantum Theory is to find a characterization of all pairs of quantum observables $\mathsf{M}_1, \mathsf{M}_2$ that satisfy $J(\mathsf{M}_1, \mathsf{M}_2) = \triangle$. Do they exist in finite dimensions? Can they have finite numbers of outcomes?

Acknowledgements

The authors wish to thank Tom Bullock for comments on an earlier version of this paper. T.H. and J.S. acknowledge financial support from the Academy of Finland (grant no. 138135).

---

[1] J. Barrett. Information processing in generalized probabilistic theories. *Phys. Rev. A*, 75:032304, 2007.

[2] G. Kimura, T. Miyadera, and H. Imai. Optimal state discrimination in general probabilistic theories. *Phys. Rev. A*, 79:062306, 2009.

[3] H. Barnum, J. Barrett, L. Orloff Clark, M. Leifer, R. Spekkens, N. Stepanik, A. Wilce, and R. Wilke. Entropy and information causality in general probabilistic theories. *New Journal of Physics*, 12:033024, 2010.

[4] G. Chiribella, G.M. D'Ariano, and P. Perinotti. Informational derivation of quantum theory. *Phys. Rev. A*, 84:012311, 2011.

[5] Paolo Perinotti. Discord and nonclassicality in probabilistic theories. *Phys. Rev. Lett.*, 108:120502, 2012.

[6] P. Busch, T. Heinonen, and P. Lahti. Heisenberg's uncertainty principle. *Phys. Rep.*, 452:155–176, 2007.

[7] P. Busch, M. Grabowski, and P.J. Lahti. *Operational Quantum Physics*. Springer-Verlag, Berlin, 1997. second corrected printing.

[8] G. Ludwig. *Foundations of Quantum Mechanics I*. Springer-Verlag, 1983.

[9] P. Busch and T. Heinosaari. Approximate joint measurements of qubit observables. *Quant. Inf. Comp.*, 8:0797–0818, 2008.

[10] P. Busch. Unsharp reality and joint measurements for spin observables. *Phys. Rev. D*, 33:2253–2261, 1986.

[11] C. Carmeli, T. Heinosaari, and A. Toigo. Informationally complete joint measurements on finite quantum systems. *Phys. Rev. A*, 85:012109, 2012.

[12] M.M. Wolf, D. Perez-Garcia, and C. Fernandez. Measurements incompatible in quantum theory cannot be measured jointly in any other no-signaling theory. *Phys. Rev. Lett.*, 103:230402, 2009.

[13] B.S. Cirel'son. Quantum generalizations of Bell's inequality. *Lett. Math. Phys.*, 4:93–100, 1980.

[14] M. Banik, M.D.R. Gazi, S. Ghosh, and G. Kar. Complementarity Principle Determines the Degree of Bell Violation in Quantum Mechanics. arXiv:1206.6054 [quant-ph], 2012.